



Southernmost records of *Hyalinobatrachium fleischmanni* (Anura: Centrolenidae)

Fernando K. Cruz,¹ Veronica L. Urgiles,^{2,4} Juan Carlos Sánchez-Nivicela,^{2,4} David C. Siddons,³ Diego F. Cisneros-Heredia^{4,5}

1 Universidad de Guayaquil, Escuela de Biología, Av. Delta y Av. Kennedy, Guayaquil, Ecuador. **2** Universidad del Azuay, Museo de Zoología, Laboratorio de Herpetología, Av. 24 de Mayo 7-77, Cuenca, Ecuador. **3** Universidad del Azuay, Escuela de Biología, Ecología y Gestión, Av. 24 de Mayo 7-77, Cuenca, Ecuador. **4** Instituto Nacional de Biodiversidad del Ecuador, División de Herpetología, calle Rumipamba 341 y Av. de los Shyris, Quito, Ecuador. **5** Universidad San Francisco de Quito USFQ, Colegio de Ciencias Biológicas y Ambientales, Laboratorio de Zoología Terrestre, Quito 170901, Ecuador.

Corresponding author: Veronica L. Urgiles, vurgiles@uazuay.edu.ec

Abstract

We present new records for Fleischmann's Glassfrog, *Hyalinobatrachium fleischmanni* (Boettger, 1893), which extend the distribution of this species into central and southwestern Ecuador and thus represent the southernmost known localities. These new reports are based on specimens collected at Cerro de Hayas, province of Guayas, and Macul, province of Los Ríos. These new localities extend the known geographic range of *H. fleischmanni* by approximately 250 km south of previous known occurrences.

Key words

Glassfrogs; western Ecuador; distribution; Guayas; Los Ríos.

Academic editor: Rodrigo Lingnau | Accepted 9 March 2017 | Accepted 30 May 2017 | Published 11 July 2017

Citation: Cruz FK, Urgiles VL, Sánchez-Nivicela JC, Siddons, DC, Cisneros-Heredia DF (2017) Southernmost records of *Hyalinobatrachium fleischmanni* (Anura: Centrolenidae). Check List 13 (4): 67–70. <https://doi.org/10.15560/13.4.67>

Introduction

Hyalinobatrachium fleischmanni (Boettger, 1893) has the widest geographic distribution of any species of the family Centrolenidae. It occurs from southern Mexico to central-western Ecuador, from sea level up to 1800 m elevation (Kubicki 2007, Frost 2016). In Ecuador, *H. fleischmanni* inhabits Lowland Seasonal and Non-Seasonal Evergreen forests in the northern and central Pacific lowlands below 800 m elevation, with records in the provinces of Esmeraldas, Manabí, Pichincha, and Los Ríos (Lynch and Duellman 1973, Cisneros-Heredia and

McDiarmid 2007a, 2007b, Delia et al. 2010).

Information regarding amphibians of western Ecuador is still fragmentary, and several species of the family Centrolenidae are known from just a few widely dispersed localities. The main explanation for this is the absence of information regarding alpha diversity for the family in the central and southern regions of Ecuador (Campos et al. 2007, Ortega-Andrade et al. 2010, Yanez-Muñoz et al. 2014). Herein, we present 2 new latitudinal records of *H. fleischmanni*, which expand its geographic range along the foothills and lowlands of central and southwestern Ecuador.



Figure 1. *Hyalinobatrachium fleischmanni*, adult male (MZUA.AN.1693) collected at Cerro de Hayas, province of Guayas, Ecuador. Photo by Fernando K. Cruz.

Methods

Specimens of *H. fleischmanni* reported herein are deposited at the Museo de Zoología, Universidad del Azuay and were collected by the authors under research permits of the Ministerio del Ambiente Ecuador (008-2015-IC-FLO/FAU-DPG/MAE and 002-2015-UB-UPN-DPALR-MAE). An adult male (Fig. 1) and a juvenile male (MZUA.AN.1693–1694) were collected at Cerro de

Hayas (−02.7299, −079.6297, 127m), approximately 20 km southwest of Naranjal, province of Guayas (Fig. 3) on 12 December 2015. One juvenile (MZUA.AN.0660, Fig. 2) and 1 adult male (MZUA.AN.0661) were collected at Macul (−01.2279 S, −079.7531, 84m), approximately 40 km southwest of Quevedo, province of Los Ríos on 16 August 2015 (Fig. 3). All the specimens were found in fragments of riparian vegetation near paddocks and pastures.

Identification

The specimens exhibit the diagnostic features described by Goin (1964), Lynch and Duellman (1973), Ortega-Andrade et al. (2010), and Wild (2003). No significant variation is distinguished among the 4 specimens, either alive or preserved. Diagnostic features include: green dorsum with small yellow spots, and yellow iris with dark spots in life; transparent parietal peritoneum; pericardium and visceral peritonea covered by iridophores (i.e. white), and all other internal peritonea transparent; extensive webbing between fingers III and IV, but basal among all other fingers; and humeral spine absent (Figs 1, 2).

Discussion

These new records significantly extend the known geographic range of *H. fleischmanni*, representing the southernmost known localities for the species. Macul is roughly 95 km south and Cerro de Hayas is approximately 250 km south from the nearest previously documented locality (Río Palenque, −00.5667, −079.3333, province of Santo Domingo; Lynch and Duellman 1973, Cisneros-



Figure 2. *Hyalinobatrachium fleischmanni*, juvenile (MZUA.AN.0660) collected at Macul, province of Los Ríos, Ecuador. Note the distinctive coloration of this species, with green dorsum and yellow spots, with a yellow iris with dark spots. Photo by Juan Carlos Sánchez-Nivicela.




Figure 3. Geographic distribution of *Hyalinobatrachium fleischmanni*. The cross represents the previous southernmost record for the species in Río Palenque (Ecuador). The triangle and square represent the new southernmost records for the species as reported herein at Macul and Cerro de Hayas (western Ecuador). Information about the distribution of *Hyalinobatrachium fleischmanni* was obtained from IUCN (2016) and Acosta-Galvis (2012).

Heredia and McDiarmid 2007b; Fig. 3). In addition, Cerro de Hayas is the first known locality of *H. fleischmanni* in the province of Guayas.

These new records reveal the presence of *H. fleischmanni* across the Pacific lowlands of Ecuador, in the Western Ecuador Biogeographic Province (sensu Morrone 2014). Despite extensive habitat destruction within this region, *H. fleischmanni* still maintains small populations in small forest fragments; more studies are needed in order to determine whether there is any need for local conservation efforts. However, as *H. fleischmanni* is a widespread species, even more studies are needed in order to explore its biogeography and taxonomic status to determine if we are indeed dealing with one species or a species complex. From this, we will be able to truly determine its local conservation status and extinction risk.

Acknowledgements

We would like to thank Ronald Navarrete, Pedro X. Astudillo, and Jacinto Guillén for institutional support. Erik Wild and Juan Manuel Guayasamin provided valuable comments to this paper. The work of VLU and JCS was

possible because of funding by Universidad del Azuay under budget code Fondos UDA 2016 [39, 2016].

Authors' Contributions

FKC, VLU and JCS collected the specimens; JCS and DFCH identified the specimens; VLU and DFCH wrote the text; VLU and JCS elaborate the map; DCS and DFCH reviewed the text and distributional maps.

References

- Acosta-Galvis AR (2012) First record of *Hyalinobatrachium fleischmanni* (Boettger, 1893) (Anura: Centrolenidae) from the Caribbean region of Colombia. Check List 8 (4): 794–795. <https://doi.org/10.15560/8.4.794>
- Boettger O (1893) Ein neuer Laubfrosch aus Costa Rica. Berichte der Senckenbergischen Naturforschenden Gesellschaft in Frankfurt am Main 1892–1893: 251–252.
- Campos F, Peralvo M, Cuesta-Camacho F, Luna S (2007) Análisis de Vacíos y Áreas Prioritarias Para la Conservación de la Biodiversidad en el Ecuador Continental. Instituto Nazca de Investigaciones Marinas / EcoCiencia / Ministerio del Ambiente / The Nature Conservancy / Conservación Internacional. Proyecto GEF: Ecuador Sistema Nacional de Áreas Protegidas, BirdLife International y Aves / Conservación, Quito, 83 pp.
- Cisneros-Heredia DF (2009) Amphibia, Anura, Centrolenidae, *Chi-*

- merella mariaelenae* (Cisneros-Heredia and McDiarmid, 2006), *Rulyrana flavopunctata* (Lynch and Duellman, 1973), *Teratohyla pulverata* (Peters, 1873), and *Teratohyla spinosa* (Taylor, 1949): historical records, distribution extension and new provincial record in Ecuador. Check List 5 (4): 912–916. <https://doi.org/10.15560/5.4.912>
- Cisneros-Heredia DF, McDiarmid RW (2005) Amphibia, Centrolenidae: *Centrolene peristictum*, *Centrolene prosoblepon*, *Cochranella cochranae*, *Cochranella midas*, *Cochranella resplendens*, *Cochranella spinosa*, *Hyalinobatrachium munozorum*: range extensions and new provincial records. Check List 1 (1): 18–22. <https://doi.org/10.15560/1.1.18>
- Cisneros-Heredia DF, McDiarmid RW (2007a) Revision of the characters of Centrolenidae (Amphibia: Anura: Athesphatanura), with comments on its taxonomy and the description of new taxa of glassfrogs. Zootaxa 1572: 1–82.
- Cisneros-Heredia DF, McDiarmid RW (2007b) Primer registro de *Hyalinobatrachium ruedai* en Ecuador con notas sobre otras especies congénéricas. Herpetotropicos 3 (1): 21–28.
- Delia J, Cisneros-Heredia DF, Whitney J, Murieta-Galindo R (2010) Observations on the reproductive behavior of a neotropical glassfrog, *Hyalinobatrachium fleischmanni* (Anura: Centrolenidae). South American Journal of Herpetology 5: 1–12. <https://doi.org/10.2994/057.005.0101>
- Frost DR (2016) Amphibian Species of the World: an Online Reference. Version 6.0. American Museum of Natural History, New York. <https://research.amnh.org/herpetology/amphibia/index.html>. Accessed on: 2016-9-13.
- Goin CJ (1964) Distribution and synonymy of *Centrolenella fleischmanni* northern South America. Herpetologica 20: 1–8.
- IUCN. 2016. The IUCN Red List of Threatened Species. Version 2016-2. International Union for Conservation of Nature. <https://www.iucnredlist.org>. Accessed on: 2016-9-13.
- Kubicki B (2007) Ranas de Vidrio Costa Rica/Glass Frogs of Costa Rica. Instituto Nacional de Biodiversidad, Santo Domingo de Heredia, 312 pp.
- Lynch JD, Duellman WE (1973) A review of the centrolenid frogs of Ecuador, with descriptions of new species. Occasional Papers of the Museum of Natural History, University of Kansas 16: 1–66.
- Morrone JJ (2014) Biogeographical regionalisation of the Neotropical region. Zootaxa 3782 (1): 1–110. <https://doi.org/10.11646/zootaxa.3782.1.1>
- Ortega-Andrade HM, Meza-Ramos P, Cisneros-Heredia DF, Yáñez-Muñoz MH, Altamirano-Benavides M (2010) Los Anfibios y Reptiles del Chocó Esmeraldeño. Serie Herpetofauna del Ecuador: El Chocó Esmeraldeño. Monografía 5. Museo Ecuatoriano de Ciencias Naturales: Quito, 95–232.
- Yáñez-Muñoz MH, Cisneros-Heredia DF (2008) Amphibia, Anura, Centrolenidae: *Cochranella orejuela*, first country records from Ecuador. Check List 4 (1): 50–54. <https://doi.org/10.15560/4.1.50>
- Yáñez-Muñoz MH, Morales M, Reyes-Puig M, Meza-Ramos P (2013) Reserva Biológica Buenaventura: Entre la transición Húmedo Tropical y la influencia Tumbesina. In: MECN, Jocotoco y Ecominga. 2013. Herpetofauna en áreas prioritarias para la conservación: El sistema de Reservas Jocotoco y Ecominga. Serie de Publicaciones del Museo Ecuatoriano de Ciencias Naturales (MECN), Monografía 6, Fundación para la Conservación Jocotoco, Fundación Ecominga. Quito, 62–75.
- Yáñez-Muñoz MH, Sánchez JC, Lopez K, Rea E, Meza-Ramos PA, Oyagata LA, Guerrero P (2014) Expansion of the distributional range of some species of amphibians and reptiles in southwestern Ecuador. Avances en Ciencias e Ingenierías 6 (1): B2–B5. <https://doi.org/10.18272/aci.v6i1.151>
- Wild ER (2003) Fleischmann's Glass Frog, *Hyalinobatrachium fleischmanni*. In: Hutchins M, Duellman WE, Schlager N (Eds) Grzimek's Animal Life Encyclopedia, 2nd edition. Vol. 6. Gale Group, Farmington Hills, Michigan, 222–223.